

Patient Name: _____ Date of Birth: _____

Health Department Use Only

Administration	TST #1	TST #2
Administrator Name:		
Date and Time Administered:		
Location:	<input type="checkbox"/> L Forearm <input type="checkbox"/> R Forearm	<input type="checkbox"/> L Forearm <input type="checkbox"/> R Forearm
Tuberculin Manufacturer:		
Tuberculin Expiration Date:		
Tuberculin Lot Number:		
Administrator Signature:		

Results (48-72 hours)	TST #1	TST #2
Date and Time Read:		
Number of mm of Induration: (<u>across</u> forearm)	_____ mm	_____ mm
Interpretation of Reading:	<input type="checkbox"/> Positive** <input type="checkbox"/> Negative	<input type="checkbox"/> Positive** <input type="checkbox"/> Negative
Reader's Signature:		

****Interpreting the TST**

≥ 5 mm is positive for:

- HIV infected
- Recent contacts
- People with fibrotic changes on CXR
- Patients with organ transplant and others on immunosuppressant drugs (including prolonged course of oral or intravenous corticosteroids or TNF alpha inhibitors)

≥ 10 mm is positive for:

- Recent immigrants (≤5 yrs) from high TB burden countries
- Injection drug users
- Mycobacterial lab workers
- People who live/work in high risk congregate settings (health care workers, long term care, correctional facilities)
- Children younger than 4 years
- Infants, children and adolescents exposed to adults in high risk categories
- People with:
Diabetes, severe kidney disease, silicosis, cancer of head or neck, hematologic or reticuloendothelial disease such as Hodgkin's disease or leukemia, intestinal bypass or gastrectomy, chronic malabsorption syndromes, low body weight

≥ 15 mm is positive if there are no known TB risk factors